



Test Definition: OPATM

Opiate Confirmation, Meconium

Overview

Useful For

Detecting maternal prenatal opiate/opioid use up to 5 months before birth

Special Instructions

- [Clinical Toxicology CPT Code Client Guidance](#)

Method Name

Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS)

NY State Available

Yes

Specimen

Specimen Type

Meconium

Ordering Guidance

For chain-of-custody testing, order OPTMX / Opiate Confirmation, Chain of Custody, Meconium.

Specimen Required

Supplies: Stool container, Small (Random), 4 oz (T288)

Container/Tube: Stool container

Specimen Volume: 1 g (approximately 1 teaspoon)

Collection Instructions: Collect entire random meconium specimen.

Forms

If not ordering electronically, complete, print, and send a [Therapeutics Test Request](#) (T831) with the specimen.

Specimen Minimum Volume

0.3 g (approximately 1/4 teaspoon)

Reject Due To

Grossly bloody	Reject; Pink OK
Stool Diapers	Reject

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Meconium	Frozen (preferred)	28 days	
	Ambient	14 days	
	Refrigerated	28 days	

Clinical & Interpretive

Clinical Information

Opiates are naturally occurring alkaloids that are derived from the opium poppy and demonstrate analgesic effects.

Opioids are derived from natural and semisynthetic alkaloids of opium or synthetic compounds(1):

-Codeine is a naturally occurring opioid agonist often incorporated into formulations along with acetaminophen or aspirin to increase its analgesic effect.(2) Codeine is metabolized to morphine and subsequently undergoes glucuronidation and sulfation.

-Morphine is an opioid receptor agonist used for major pain analgesia.(2) It has been shown to distribute widely into many fetal tissues(3) and has been detected in meconium.

-Hydrocodone is a semisynthetic analgesic derived from codeine. Hydrocodone is 6 times more potent than codeine and is prescribed for treatment of moderate-to-moderately severe pain.(2) Hydrocodone undergoes O-demethylation in vivo, forming hydromorphone.

-Hydromorphone, a semisynthetic derivative of morphine, is an opioid analgesic. It is 7 to 10 times more potent than morphine, its addiction liability is similar to morphine.(2)

-Oxycodone, a semisynthetic narcotic derived from thebaine. It is metabolized by O-demethylation, forming oxymorphone.(2)

-Oxymorphone is a semisynthetic opioid derivative of thebaine and is indicated for moderate-to-severe pain.(2)

-Heroin, a semisynthetic derivative of morphine, is rapidly deacetylated in vivo to the active metabolite 6-monoacetylmorphine (6-MAM), which is further hydrolyzed to morphine.(2)

Opiates have been shown to readily cross the placenta and distribute widely into many fetal tissues. Opiate use by the mother during pregnancy increases the risk of prematurity and small size for gestational age. Furthermore, heroin-exposed infants exhibit an early onset of withdrawal symptoms compared to methadone-exposed infants. These infants demonstrate a variety of symptoms, including irritability, hypertonia, wakefulness, diarrhea, yawning, sneezing, increased hiccups, jitteriness, excessive sucking, and seizures. Long-term intrauterine drug exposure may lead to abnormal neurocognitive and behavioral development as well as an increased risk of sudden infant death syndrome.

The disposition of opiates and opioids in meconium, the first fecal material passed by the neonate, is not well understood. The proposed mechanism is that the fetus excretes drug into bile and amniotic fluid. Drug accumulates in meconium either by direct deposition from bile or through swallowing of amniotic fluid. The first evidence of meconium in the fetal intestine appears at approximately the 10th to 12th week of gestation, and it slowly moves into the colon by the 16th week of gestation. Therefore, the presence of drugs in meconium has been proposed to be indicative of in utero drug exposure during the final 4 to 5 months of pregnancy, a longer historical measure than is possible by urinalysis.

Reference Values

Negative

Positives are reported with a quantitative liquid chromatography tandem mass spectrometry (LC-MS/MS) result.

Cutoff concentrations for LC-MS/MS testing:

Codeine: 20 ng/g

Hydrocodone: 20 ng/g

Hydromorphone: 20 ng/g

Morphine: 20 ng/g

Oxycodone: 20 ng/g

Oxymorphone: 20 ng/g

Interpretation

The presence of any of the following opiates (codeine, morphine, hydrocodone, hydromorphone, oxycodone, oxymorphone) at 20 ng/g or greater or 6-monoacetylmorphine at 10 ng/g or greater indicates the newborn was exposed to opiates/opioids during gestation.

Cautions

No significant cautionary statements

Clinical Reference

1. Gutstein HB, Akil H. Opioid analgesics. In: Brunton LL, Lazo JS, Parker KL, eds. Goodman and Gilman's The Pharmacological Basis of Therapeutics. 11th ed. McGraw-Hill; 2006
2. Baselt RC. Disposition of Toxic Drugs and Chemical in Man. 12th ed. Biomedical Publications; 2020
3. Szeto HH. Kinetics of drug transfer to the fetus. Clin Obstet Gynecol. 1993;36(2):246-254
4. Ahanya SN, Lakshmanan J, Morgan BLG, Ross MG. Meconium passage in utero: mechanisms, consequences, and management. Obstet Gynecol Surv. 2005;60(1):45-56
5. Langman LJ, Bechtel LK, Holstege CP. Clinical toxicology. In: Rifai N, Chiu RWK, Young I, Burnham C-AD, Wittwer CT, eds. Tietz Textbook of Laboratory Medicine. 7th ed. Elsevier; 2023:454-454: chap 43

Performance

Method Description

Meconium is mixed with internal standard and extracted with methanol. The methanolic extract is further processed by solid phase extraction. The extract is analyzed by liquid chromatography tandem mass spectroscopy.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Sunday

Report Available

2 to 3 days

Specimen Retention Time

2 weeks

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Superior Drive

Fees & Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

G0480

80361 (if appropriate for select payers)

80365 (if appropriate for select payers)

[Clinical Toxicology CPT Code Client Guidance](#)**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
OPATM	Opiate Confirmation, M	69026-3

Result ID	Test Result Name	Result LOINC® Value
31848	Morphine	69027-1
31850	Oxymorphone	69028-9
31849	Hydromorphone	68541-2
31847	Codeine	68542-0
31852	Oxycodone	68543-8
31851	Hydrocodone	68544-6
31868	Interpretation	8215-6
31869	Chain of Custody	77202-0