

Overview

Useful For

Detecting mercury toxicity in 24-hour urine specimens

Special Instructions

- [Urine Preservatives-Collection and Transportation for 24-Hour Urine Specimens](#)
- [Metals Analysis Specimen Collection and Transport](#)

Method Name

Triple Quadrupole Inductively Coupled Plasma-Mass Spectrometry (ICP-MS/MS)

NY State Available

Yes

Specimen

Specimen Type

Urine

Necessary Information

24-Hour volume (in milliliters) is required.

Specimen Required

Patient Preparation: High concentrations of gadolinium and iodine are known to interfere with most metal tests. If either gadolinium- or iodine-containing contrast media has been administered, a specimen should not be collected for 96 hours.

Supplies: Urine Tubes, 10 mL (T068)

Collection Container/Tube: Clean, plastic urine container with no metal cap or glued insert

Submission Container/Tube: Plastic, 10-mL urine tube or a clean, plastic aliquot container with no metal cap or glued insert

Specimen Volume: 10 mL

Collection Instructions:

1. Collect urine for 24 hours.
2. Refrigerate specimen within 4 hours of completion of 24-hour collection.
3. See [Metals Analysis Specimen Collection and Transport](#) for complete instructions.

Additional Information: See [Urine Preservatives-Collection and Transportation for 24-Hour Urine Specimens](#) for multiple collections.

Urine Preservative Collection Options

Note: The addition of preservative or application of temperature controls **must occur within 4 hours of completion of**

the collection.

Ambient	No
Refrigerate	Preferred
Frozen	OK
50% Acetic Acid	OK
Boric Acid	No
Diazolidinyl Urea	No
6M Hydrochloric Acid	OK
6M Nitric Acid	OK
Sodium Carbonate	No
Thymol	No
Toluene	No

Specimen Minimum Volume

2 mL

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	7 days	
	Frozen	7 days	

Clinical & Interpretive

Clinical Information

The correlation between the levels of mercury (Hg) excretion in the urine and the clinical symptoms is considered poor.

Previous thought indicated urine as a more appropriate marker of inorganic mercury because organic mercury represented only a small fraction of urinary mercury. Based on possible demethylation of methylmercury within the body, urine may represent a mixture of dietary methylmercury and inorganic mercury. Seafood consumption can contribute to urinary mercury levels (up to 30%),⁽¹⁾ which is consistent with the suggestion that due to demethylation processes in the human body, a certain proportion of urinary mercury can originate from dietary consumption of fish/seafood.⁽²⁾

For more information see HG / Mercury, Blood.

Reference Values

- 0-17 years: Not established
- > or =18 years: <2 mcg/24 h
- Toxic concentration: >50 mcg/24 h

The concentration at which toxicity is expressed is widely variable between patients. 50 mcg/24 h is the lowest concentration at which toxicity is usually apparent.

Interpretation

Daily urine excretion of mercury above 50 mcg/day indicates significant exposure (per World Health Organization standard).

Cautions

To avoid contamination by dust, specimen should be collected away from the site of suspected exposure.

Clinical Reference

1. Snoj Tratniid J, Falnoga I, Mazej D, et al: Results of the first national human biomonitoring in Slovenia: Trace elements in men and lactating women, predictors of exposure and reference values. *Int J Hyg Environ Health*. 2019 Apr;222(3):563-582
2. Sherman LS, Blum JD, Franzblau A, Basu N: New insights into biomarkers of human mercury exposure using naturally occurring mercury stable isotopes. *Environ Sci Technol*. 2013 Apr 2;47(7):3403-3409
3. Lee R, Middleton D, Caldwell K, et al: A review of events that expose children to elemental mercury in the United States. *Environ Health Perspect*. 2009 Jun;117(6):871-878
4. Bjorkman L, Lundekvam BF, Laegreid T, et al: Mercury in human brain, blood, muscle and toenails in relation to exposure: an autopsy study. *Environ Health*. 2007 Oct 11;6:30
5. Bernhoft RA. Mercury toxicity and treatment: a review of the literature. *J Environ Public Health*. 2012;2012:460508. doi: 10.1155/2012/460508
6. Strathmann FG, Blum LM: Toxic elements. In: Rifai N, Chiu RWK, Young I, Burnham CD, Wittwer CT, eds. *Tietz Textbook of Laboratory Medicine*. 7th ed. Elsevier; 2023:chap 44

Performance**Method Description**

The metal of interest is analyzed by inductively coupled plasma mass spectrometry.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 3 days

Specimen Retention Time

14 days

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

83825

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
HGU	Mercury, 24 Hr, U	6693-6

Result ID	Test Result Name	Result LOINC® Value
8592	Mercury, 24 Hr, U	6693-6
TM5	Collection Duration	13362-9
VL3	Urine Volume	3167-4