

Test Definition: FSOMA

Somatostatin (Somatotropin Release-Inhibiting Factor, SRIF)

Overview

Method Name

Direct Radioimmunoassay (RIA)

NY State Available

No

Specimen

Specimen Type

GI Plasma

Specimen Required

Patient preparation:

1. Patient should be fasting 10-12 hours prior to collection.

2. Patient should not be on any medications that affect insulin secretion or intestinal motility, if possible for at least 48 hours prior to collection.

Specimen Type: GI Plasma

Container/Tube: EDTA tube containing GI preservative (T125)

Specimen Volume: 3 mL

Collection Instructions: Collect 10 mL of blood in special tube containing G.I. Preservative (T125). Specimen should be separated immediately and send 3 mL plasma frozen as soon as possible.

Specimen Minimum Volume

1 mL

Reject Due To

Gross	Reject
hemolysis	
Gross lipemia	Reject
Gross icterus	Reject
Other	Specimens other than Plasma collected in GI preservative (T125).

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
GI Plasma	Frozen	180 days	



Test Definition: FSOMA

Somatostatin (Somatotropin Release-Inhibiting Factor, SRIF)

Clinical & Interpretive

Clinical Information

Somatostatin is a cyclic peptide originally isolated from sheep hypothalami and shown to inhibit the release of Growth Hormone. Somatostatin is present primarily in three main forms: a 14 amino acid peptide, a 28 amino acid peptide ("Big" Somatostatin), and a 12,000 molecular weight ProSomatostatin. This assay measures only the 14 amino acid form of Somatostatin. All three forms of Somatostatin have similar biological properties and overall potencies. Somatostatin is a physiological regulator of islet cell and gastrointestinal functions and is a suppressor of many pituitary hormones including Growth Hormone, Prolactin, and Thyrotropin (TSH). Somatostatin levels are often elevated in diabetics, but the levels return to normal upon correction of the hormonal and metabolic deficiencies present. In many cases of APUDomas including VIPoma, Insulinoma, Glucagonoma, and Gastrinoma, elevated levels of Somatostatin are found.

Reference Values

Up to 25 pg/mL

Performance

PDF Report

Day(s) Performed Monday through Friday

Report Available 9 to 11 days

Performing Laboratory Location

Inter Science Institute

Fees & Codes

Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact Customer Service.

Test Classification

This test was developed and its performance characteristics determined by Inter Science Institute. It has not been cleared or approved by the US Food and Drug Administration. The FDA has determined that such clearance or approval



Test Definition: FSOMA

Somatostatin (Somatotropin Release-Inhibiting Factor, SRIF)

is not necessary.

CPT Code Information

84307

LOINC[®] Information

Test ID	Test Order Name	Order LOINC [®] Value
FSOMA	Somatostatin (Plasma)	2961-1
Result ID	Test Result Name	Result LOINC [®] Value
Z0933	Somatostatin, (Plasma)	2961-1