

Overview

Useful For
Aiding in the diagnosis of neurosyphilis

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
VDSFQ	VDRL Titer, CSF	No	No

Testing Algorithm
If this test is positive, a VDRL titer will be performed at an additional charge.

For more information, see [Meningitis/Encephalitis Panel Algorithm](#).

Special Instructions
• [Meningitis/Encephalitis Panel Algorithm](#)

Method Name
Flocculation/Agglutination

NY State Available
Yes

Specimen

Specimen Type
CSF

Specimen Required
Collection Container/Tube: Sterile vial
Submission Container/Tube: Plastic vial
Specimen Volume: 0.5 mL
Collection Instructions: Submit specimen collected in vial 2, if possible. If not, note which vial from which the aliquot was obtained.

Specimen Minimum Volume
0.2 mL

Reject Due To

Gross hemolysis	Reject
Gross lipemia	Reject

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
CSF	Frozen (preferred)	14 days	
	Refrigerated	14 days	

Clinical & Interpretive

Clinical Information

VDRL is a nontreponemal serologic test for syphilis that uses a cardiolipin-cholesterol-lecithin antigen to detect reaginic antibodies. The VDRL test performed on cerebrospinal fluid can be used to diagnose neurosyphilis in patients with a prior history of syphilis infection.

The presence of neurosyphilis in untreated patients can be detected by the presence of pleocytosis, elevated protein, and a positive VDRL.

Reference Values

Negative
Reference values apply to all ages.

Interpretation

A positive VDRL result on spinal fluid is highly specific for neurosyphilis.

A single negative VDRL result should not be used to exclude neurosyphilis and repeat testing on a new specimen may be necessary.

Positive results will be titered.

Cautions

VDRL testing on spinal fluid gives a high percentage of false-negative results.

Clinical Reference

1. Miller JN: Value and limitations of nontreponemal and treponemal tests in the laboratory diagnosis of syphilis. Clin Obstet Gynecol. 1975 Mar;18(1):191-203

2. Radolf JD, Tramont EC, Salazar JC: Syphilis (*Treponema pallidum*). In: Bennett JE, Dolin R, Blaser MJ, eds. Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases. 9th ed. Elsevier; 2020:2865-2892

Performance

Method Description

VDRL antigen and spinal fluid are mixed on a 180 RPM rotator. The antigen, a cardiolipin-lecithin coated cholesterol particle, flocculates in the presence of reagin.(US Department of Health, Education and Welfare, National Communicable Diseases Center, Venereal Disease Program: Manual of Tests for Syphilis. Centers for Disease Control; 1969; Bennett JE, Dolin R, Blaser MJ, eds. Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases. 9th ed. Elsevier; 2020)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 4 days

Specimen Retention Time

14 days

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

86592

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
VDSF	VDRL, CSF	5290-2

Result ID	Test Result Name	Result LOINC® Value
9028	VDRL, CSF	5290-2