



# Test Definition: FHIPP

Toluene as Hippuric Acid, Occupational Exposure, Urine

## Overview

### Method Name

Gas Chromatography/Flame Ionization Detection (GC-FID)

### NY State Available

Yes

## Specimen

### Specimen Type

Urine

### Specimen Required

10 mL aliquot of random or spot urine collected at end of shift. Send specimen refrigerated. Tubes should be filled to prevent loss of volatile compound into headspace.

### Specimen Minimum Volume

1.2 mL

### Reject Due To

Hemolysis	NA
Lipemia	NA
Icterus	NA
Other	NA

### Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	14 days	
	Frozen	180 days	

## Clinical & Interpretive

### Reference Values

Creatinine: >50 mg/cL

---

Hippuric Acid is a metabolite of toluene and benzyl alcohol.

Normal (unexposed population):

Average 0.8 g/L

Exposed:

Biological Exposure Index (BEI):

1.6 g/g creatinine

(toluene exposure: end of shift)

Toxic:

Not established

## Performance

### PDF Report

No

### Day(s) Performed

Monday through Sunday

### Report Available

7 to 8 days

### Specimen Retention Time

2 weeks

### Performing Laboratory Location

Medtox Laboratories, Inc.

## Fees & Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

### CPT Code Information

83921

82570

---

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
FHIPP	Toluene as Hippuric Acid, Occup, U	Not Provided

Result ID	Test Result Name	Result LOINC® Value
Z1863	Creatinine	2161-8
Z1060	Hippuric Acid	6709-0
Z1061	Hippuric Acid	13754-7