

Test Definition: TAMIK

Amikacin, Trough, Serum

Reporting Title: Amikacin, Trough, S

Performing Location: Rochester

Specimen Requirements:

Collection Container/Tube: Preferred: Serum gel Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 0.5 mL Collection Instructions:

1. Serum gel tubes should be centrifuged within 2 hours of collection.

2. Red-top tubes should be centrifuged, and the serum aliquoted into a plastic vial within 2 hours of collection.

Specimen Minimum Volume:

0.25 mL

Forms:

If not ordering electronically, complete, print, and send a Therapeutics Test Request (T831) with the specimen.

| Specimen Type | Temperature | Time | Special Container |
|---------------|--------------------------|----------|-------------------|
| Serum | Refrigerated (preferred) | 14 days | |
| | Frozen | 28 days | |
| | Ambient | 72 hours | |

Result Codes:

| Result ID | Reporting Name | Туре | Unit | LOINC® |
|-----------|---------------------|---------|--------|--------|
| TAMIK | Amikacin, Trough, S | Numeric | mcg/mL | 3321-7 |

LOINC and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

80150



Test Definition: TAMIK

Amikacin, Trough, Serum

Reference Values:

Trough: <8.0 mcg/mL Toxic trough: >10.0 mcg/mL