

Test Definition: TOBPA

Tobramycin, Peak, Serum

Reporting Title: Tobramycin, Peak, S

Performing Location: Rochester

Specimen Requirements:

Collection Container/Tube: Preferred: Serum gel Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 0.5 mL Collection Instructions:

1. Serum gel tubes should be centrifuged within 2 hours of collection.

2. Red-top tubes should be centrifuged, and the serum aliquoted into a plastic vial within 2 hours of collection.

Specimen Minimum Volume:

0.25 mL

Forms:

If not ordering electronically, complete, print, and send a Therapeutics Test Request (T831) with the specimen.

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	7 days	
	Frozen	28 days	
	Ambient	72 hours	

Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
ТОВРА	Tobramycin, Peak, S	Numeric	mcg/mL	4057-6

LOINC and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

80200



Test Definition: TOBPA

Tobramycin, Peak, Serum

Reference Values:

Therapeutic: 3.0-12.0 mcg/mL Toxic: >12.0 mcg/mL