

**Reporting Title:** Activated Partial Thromboplastin Time, P  
**Performing Location:** Rochester

**Necessary Information:**

Heparin or Coumadin therapy should be noted.

**Specimen Requirements:**

Only orderable as part of a special coagulation profile or as a reflex. For more information see:

ALUPP / Lupus Anticoagulant Profile, Plasma

ALBLD / Bleeding Diathesis Profile, Limited, Plasma

AATHR / Thrombophilia Profile, Plasma and Whole Blood

APROL / Prolonged Clot Time Profile, Plasma

ADIC / Disseminated Intravascular Coagulation/Intravascular Coagulation and Fibrinolysis (DIC/ICF) Profile, Plasma

Specimen Type	Temperature	Time	Special Container
Plasma Na Cit	Frozen	14 days	

**Result Codes:**

Result ID	Reporting Name	Type	Unit	LOINC®
APTSC	Activated Partial Thromboplastin Time, P	Numeric	sec	14979-9

LOINC and CPT codes are provided by the performing laboratory.

**Supplemental Report:**

No

**CPT Code Information:**

85730

**Reference Values:**

Only orderable as part of a special coagulation profile or as a reflex. For more information see:

ALUPP / Lupus Anticoagulant Profile, Plasma

ALBLD / Bleeding Diathesis Profile, Limited, Plasma

AATHR / Thrombophilia Profile, Plasma and Whole Blood

APROL / Prolonged Clot Time Profile, Plasma

ADIC / Disseminated Intravascular Coagulation/Intravascular Coagulation and Fibrinolysis (DIC/ICF) Profile, Plasma

---

25-37 seconds

The APTT may be 35% longer in full-term newborns that reach adult reference range by age 3 months and twice the adult upper limit in premature infants reaching adult reference range by age 6 months.