

Test Definition: CRCRU

Chromium/Creatinine Ratio, Random, Urine

Reporting Title: Chromium/Creat Ratio, Random, U **Performing Location:** Rochester

Ordering Guidance:

High concentrations of gadolinium and iodine are known to interfere with most metal tests. If either gadolinium- or iodine-containing contrast media has been administered, a specimen should not be collected for 96 hours.

Specimen Requirements:

Supplies: Urine Tubes, 10 mL (T068) Collection Container/Tube: Clean, plastic urine container with no metal cap or glued insert Submission Container/Tube: Plastic vial or clean, plastic aliquot container with no metal cap or glued insert Specimen Volume: 3 mL

Collection Instructions:

1. Collect a random urine specimen.

2. See Metals Analysis Specimen Collection and Transport for complete instructions.

Specimen Minimum Volume:

2 mL

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	28 days	
	Frozen	28 days	
	Ambient	14 days	

Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
607759	Chromium/Creat Ratio, U	Numeric	mcg/g Cr	13464-3
CRETR	Creatinine, Random, U	Numeric	mg/dL	2161-8

LOINC and CPT codes are provided by the performing laboratory.

Supplemental Report:

No



Chromium/Creatinine Ratio, Random, Urine

Components:

Test ID	Reporting Name	CPT Units	CPT Code	Always Performed	Orderable Separately
CRCRR	Chromium/Creat Ratio, U			Yes	No
CRETR	Creatinine, Random, U			Yes	No

CPT Code Information:

82495 82570

Reference Values:

0-17 years: Not established

>17 years: <0.8 mcg/g Creatinine