

**Reporting Title:** Calcitonin, FNAB, Lymph Node **Performing Location:** Rochester

### Shipping Instructions:

Send specimen frozen.

### **Necessary Information:**

The biopsied site of each specimen is required and must be clearly identified in the LIS and/or batch sheet.

### Specimen Requirements:

Patient Preparation: For 12 hours before this procedure do not take multivitamins or dietary supplements containing biotin (vitamin B7), which is commonly found in hair, skin, and nail supplements and multivitamins.

Container/Tube: Plain, plastic, screw-top tube

Specimen Volume: 1 to 1.5 mL

Collection Instructions:

1. Needle wash specimens for analysis should be collected in conjunction with cytology specimens.

2. Have saline available prior to start of procedure. Saline is the only acceptable solution for needle washings.

3. After each fine-needle aspiration biopsy (FNAB) has been collected and the material in the needle has been expelled onto a slide for cytologic analysis, attach the used FNAB needle to an empty syringe.

4. Withdraw between 0.10 mL and 0.25 mL of saline up through the needle until the saline starts to fill the hub of the needle or end of the syringe.

5. Expel this fluid back through the needle into a separate tube. This is the needle washing used for analysis.

6. Repeat steps 2 through 4 for each needle pass of the same biopsied site and empty into the same tube, accumulating a total of 0.5 mL to 1.5 mL of fluid to send to the laboratory. (If more than 1 site is biopsied, see Additional Information)

7. Inspect specimen for visible blood or tissue contamination:

a. If bloody, centrifuge specimen and transfer supernatant to a new plastic aliquot tube (5-mL standard tube) to send to laboratory. The supernatant, not the cellular material, is used for analysis.

b. If specimen is clear, centrifugation is not necessary.

8. Freeze within 2 to 4 hours of collection.

Additional Information:

1. If more than 1 site is biopsied, each washing material should be submitted on a separate tube and under a different order number.

2. A minimum of 0.5 mL is required for testing; however, the total collection volume should not exceed 1.5 mL. Sample volumes outside these parameters may be rejected.

3. Do not send saline control. This test has been validated to rule-out saline matrix effect.

### Specimen Minimum Volume:

0.5 mL

# Forms:

If not ordering electronically, complete, print, and send an Oncology Test Request (T729) with the specimen.



Calcitonin, Fine-Needle Aspiration Biopsy Needle Wash, Lymph Node

Specimen Type	Temperature	Time	Special Container
Fine Needle Wash	Frozen (preferred)	7 days	
	Refrigerated	4 hours	

# Ask at Order Entry (AOE) Questions:

Test ID	Question ID	Description	Туре	Reportable
CATLN	SITEB	Site	Plain Text	Yes

# Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
CATL	Calcitonin, FNAB, Lymph Node	Numeric	pg/mL	75709-6
SITEB	Site	Alphanumeric		39111-0

LOINC and CPT codes are provided by the performing laboratory.

### Supplemental Report:

No

# **CPT Code Information:**

82308

# **Reference Values:**

An interpretive report will be provided.