

Antimony, Blood

## Reporting Title: Antimony, B

Performing Location: Rochester

#### Specimen Requirements:

Patient Preparation: High concentrations of gadolinium and iodine are known to interfere with most metal tests. If either gadolinium- or iodine-containing contrast media has been administered, a specimen should not be collected for 96 hours.

Supplies: Greiner Bio-One VACUETTE TUBE 6 mL NH Trace Elements Sodium Heparin tube (T819)

Container/Tube: Greiner Bio-One Vacuette Tube 6 mL NH Trace Elements Sodium Heparin tube for blood is required. Becton-Dickinson (BD) royal blue-top tubes are not acceptable for this testing.

Specimen Volume: 0.8 mL

Collection Instructions:

- 1. See Metals Analysis Specimen Collection and Transport for complete instructions.
- 2. Send whole blood specimen in original tube. Do not aliquot.

#### **Specimen Minimum Volume:**

0.25 mL

Specimen Type	Temperature	Time	Special Container
Whole blood	Refrigerated	7 days	

## **Result Codes:**

Result ID	Reporting Name	Туре	Unit	LOINC®
64273	Antimony, B	Numeric	ng/mL	5578-0

LOINC and CPT codes are provided by the performing laboratory.

## Supplemental Report:

No

## **CPT Code Information:**

83018



# **Test Definition: SBWB**

Antimony, Blood

#### **Reference Values:**

<3 ng/mL (unexposed) 3-10 ng/mL (exposed)