

Test Definition: PSPT

Phosphatidylserine/Prothrombin Antibody, IgG and IgM, Serum

Reporting Title: PS/PT Ab, IgG/IgM, S

Performing Location: Rochester

Ordering Guidance:

Cardiolipin and beta-2 glycoprotein testing are the first-tier test options for most patients. Phosphatidylserine/prothrombin antibodies are considered part of the second-tier workup.

Specimen Requirements:

Supplies: Sarstedt Aliquot Tube, 5 mL (T914)

Collection Container/Tube: Preferred: Serum gel Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 0.5 mL

Collection Instructions: Centrifuge and aliquot serum into a plastic vial.

Specimen Minimum Volume:

0.4 mL

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	21 days	
	Frozen	21 days	

Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
PSPTG	PS/PT Ab, IgG, S	Numeric	U	85359-8
	Also used by tests: PSPTG			
PSPTM	PS/PT Ab, IgM, S	Numeric	U	85358-0
	Also used by tests: PSPTM			

LOINC and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

Test Definition: PSPT

Phosphatidylserine/Prothrombin Antibody, IgG and IgM, Serum

Components:

Test ID	Reporting Name	CPT Units	CPT Code	Always Performed	Orderable Separately
PSPTG	PS/PT Ab, IgG, S			Yes	Yes
PSPTM	PS/PT Ab, IgM, S			Yes	Yes

CPT Code Information:

86148 x 2

Reference Values:

Negative < or =30.0 U Borderline 30.1-40.0 U Positive > or =40.1 U