

Test Definition: CH9

Chromogenic Factor IX Activity Assay, Plasma

Reporting Title: Chromogenic FIX, P

Performing Location: Rochester

Ordering Guidance:

Coagulation testing is highly complex, often requiring the performance of multiple assays and correlation with clinical information. For that reason, ALBLD / Bleeding Diathesis Profile, Limited, Plasma is recommended.

Specimen Requirements:

Specimen Type: Platelet-poor plasma

Collection Container/Tube: Light-blue top (3.2% sodium citrate) Submission Container/Tube: Polypropylene vial preferred

Specimen Volume: 1 mL Collection Instructions:

- 1. For complete instructions, see Coagulation Guidelines for Specimen Handling and Processing
- 2. Centrifuge, transfer all plasma into a plastic vial, and centrifuge plasma again.
- 3. Aliquot plasma into a separate plastic vial leaving 0.25 mL in the bottom of centrifuged vial.
- 4. Freeze plasma immediately (no longer than 4 hours after collection) at -20 degrees C or, ideally, at -40 degrees C or below.

Additional Information:

- 1. Double-centrifuged specimen is critical for accurate results as platelet contamination may cause spurious results.
- 2. Each coagulation assay requested should have its own vial.

Specimen Minimum Volume:

0.5 mL

Forms:

If not ordering electronically, complete, print, and send a Coagulation Test Request (T753) with the specimen.

Specimen Type	Temperature	Time	Special Container
Plasma Na Cit	Frozen	14 days	

Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
CH9	Chromogenic FIX, P	Numeric	%	88449-4

LOINC and CPT codes are provided by the performing laboratory.



Test Definition: CH9

Chromogenic Factor IX Activity Assay, Plasma

Supplemental	Report:
--------------	---------

No

CPT Code Information:

85130

Reference Values:

65-140%

Chromogenic factor IX activity generally correlates with the one-stage FIX activity. In full term/premature neonates, infants, children, and adolescents the one-stage FIX activity* is similar to adults. However, no similar data for chromogenic FIX activity are available.(Appel IM, Grimminck B, Geerts J, Stigter R, Cnossen MH, Beishuizen A. Age dependency of coagulation parameters during childhood and puberty. J Thromb Haemost. 2012;10(11):2254-2263)

*See Pediatric Hemostasis References section in Coagulation Guidelines for Specimen Handling and Processing