

**Reporting Title:** Aldosterone, IVC**Performing Location:** Rochester**Specimen Requirements:**

Patient Preparation: Spironolactone (Aldactone) should be discontinued for 4 to 6 weeks before testing. The plasma renin activity cannot be interpreted if the patient is being treated with spironolactone.

Collection Container/Tube:

Preferred: Red top

Acceptable: Serum gel

Submission Container/Tube: Plastic vial

Specimen Volume: 1.8 mL

Collection Instructions:

1. See Renin-Aldosterone Studies for detailed instructions.
2. The recommended collection time is 8 a.m., after the patient is active for approximately 2 hours. Try to collect the specimen as close to that time as possible and no later than 10 a.m.
3. Centrifuge and aliquot serum into a plastic vial.

**Specimen Minimum Volume:**

1 mL

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	28 days	
	Frozen	30 days	
	Ambient	4 days	

**Result Codes:**

Result ID	Reporting Name	Type	Unit	LOINC®
6503	Aldosterone, IVC	Numeric	ng/dL	1763-2

LOINC and CPT codes are provided by the performing laboratory.

**Supplemental Report:**

No

**CPT Code Information:**

82088

---

**Reference Values:**

No established reference values.