

Test Definition: PFN

Propafenone, Serum

Reporting Title: Propasenone, S **Performing Location:** Rochester

Specimen Requirements:

Patient Preparation: Specimens should only be collected after patient has been receiving propafenone for at least 3

days. Trough concentrations should be collected just before administration of the next dose.

Collection Container/Tube: Red top (serum gel/SST are not acceptable)

Submission Container/Tube: Plastic vial

Specimen Volume: 1.5 mL Collection Instructions:

1. Draw blood immediately before next scheduled dose.

2. Centrifuge and aliquot serum into plastic vial within 2 hours of collection.

Specimen Minimum Volume:

0.5 mL

Forms:

If not ordering electronically, complete, print, and send a Therapeutics Test Request (T831) with the specimen.

| Specimen Type | Temperature | Time | Special Container |
|---------------|--------------------------|---------|-------------------|
| Serum Red | Refrigerated (preferred) | 28 days | |
| | Ambient | 28 days | |
| | Frozen | 28 days | |

Result Codes:

| Result ID | Reporting Name | Туре | Unit | LOINC® |
|-----------|----------------|---------|--------|--------|
| 80295 | Propafenone, S | Numeric | mcg/mL | 6905-4 |

LOINC and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

80299



Test Definition: PFN

Propafenone, Serum

Reference Values:

Trough Value

0.5-2.0 mcg/mL: Therapeutic concentration

>2.0 mcg/mL: Toxic concentration