

Inhibin A, Tumor Marker, Serum

# **Reporting Title:** Inhibin A, Tumor Marker, S **Performing Location:** Rochester

#### Ordering Guidance:

For the initial evaluation of patients suspected of having a granulosa cell tumor of the ovary, order INHAB / Inhibin A and B, Tumor Marker, Serum. If the results of the profile show that either inhibin A or B are elevated, consider monitoring the patient with the individual tests, INHA / Inhibin A, Tumor Marker, Serum or INHB / Inhibin B, Serum.

#### Specimen Requirements:

Supplies: Sarstedt Aliquot Tube 5 mL (T914) Collection Container/Tube: Preferred: Serum gel Acceptable: Red top Submission Container/Tube: Plastic vial Specimen Volume: 0.6 mL Collection Instructions: Centrifuge and aliquot serum into a plastic vial.

#### **Specimen Minimum Volume:**

0.5 mL

#### Forms:

If not ordering electronically, complete, print, and send an Oncology Test Request (T729) with the specimen.

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	14 days	
	Frozen	90 days	
	Ambient	7 days	

#### Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
INHA	Inhibin A, Tumor Marker, S	Numeric	pg/mL	23883-2

LOINC and CPT codes are provided by the performing laboratory.

#### Supplemental Report:

No



# **Test Definition: INHA**

Inhibin A, Tumor Marker, Serum

## **CPT Code Information:**

86336

### **Reference Values:**

Males: <5.0 pg/mL

Females <11 years: <5.0 pg/mL 11-17 years: <98 pg/mL Premenopausal: <98 pg/mL Postmenopausal: <5.0 pg/mL