

Reporting Title: Testosterone, Total, Bio, Free, S**Performing Location:** Rochester**Ordering Guidance:**

The preferred test for diagnosis of mild abnormalities of testosterone homeostasis, particularly if abnormalities in sex hormone-binding globulin function or levels are present, is TTBS / Testosterone, Total and Bioavailable, Serum.

Necessary Information:

Patient's age and sex are required.

Specimen Requirements:

Supplies: Sarstedt Aliquot Tube, 5 mL (T914)

Collection Container/Tube: Red top (serum gel/SST are not acceptable)

Submission Container/Tube: Plastic vial

Specimen Volume: 3.5 mL

Collection Instructions: Centrifuge and aliquot serum into a plastic vial

Specimen Minimum Volume:

2 mL

Specimen Type	Temperature	Time	Special Container
Serum Red	Refrigerated (preferred)	14 days	
	Frozen	60 days	

Result Codes:

Result ID	Reporting Name	Type	Unit	LOINC®
8533	Testosterone, Total, S Also used by tests: TTST	Numeric	ng/dL	2986-8
3631	Testosterone Free	Numeric	ng/dL	2991-8
82978	Testosterone, Bioavailable, S	Numeric	ng/dL	2990-0

LOINC and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

Components:

Test ID	Reporting Name	CPT Units	CPT Code	Always Performed	Orderable Separately
TTST	Testosterone, Total, S			Yes	Yes
FRTST	Testosterone, Free, S			Yes	No
BATS	Testosterone, Bioavailable, S			Yes	No

CPT Code Information:

84402
84403
84410

Reference Values:**TESTOSTERONE, TOTAL****Males**

0-5 months: 75-400 ng/dL
6 months-9 years: <7-20 ng/dL
10-11 years: <7-130 ng/dL
12-13 years: <7-800 ng/dL
14 years: <7-1,200 ng/dL
15-16 years: 100-1,200 ng/dL
17-18 years: 300-1,200 ng/dL
> or =19 years: 240-950 ng/dL
Tanner Stages*
I (prepubertal): <7-20
II: 8-66
III: 26-800
IV: 85-1,200
V (young adult): 300-950

Females

0-5 months: 20-80 ng/dL
6 months-9 years: <7-20 ng/dL
10-11 years: <7-44 ng/dL
12-16 years: <7-75 ng/dL
17-18 years: 20-75 ng/dL
> or =19 years: 8-60 ng/dL
Tanner Stages*
I (prepubertal): <7-20
II: <7-47
III: 17-75
IV: 20-75

V (young adult): 12-60

*Puberty onset (transition from Tanner stage I to Tanner stage II) occurs for boys at a median age of 11.5 (+/-2) years and for girls at a median age of 10.5 (+/-2) years. There is evidence that it may occur up to 1 year earlier in obese girls and in African American girls. For boys, there is no definite proven relationship between puberty onset and body weight or ethnic origin. Progression through Tanner stages is variable. Tanner stage V (young adult) should be reached by age 18.

TESTOSTERONE, FREE

Males (adult):

20-<25 years: 5.25-20.7 ng/dL
25-<30 years: 5.05-19.8 ng/dL
30-<35 years: 4.85-19.0 ng/dL
35-<40 years: 4.65-18.1 ng/dL
40-<45 years: 4.46-17.1 ng/dL
45-<50 years: 4.26-16.4 ng/dL
50-<55 years: 4.06-15.6 ng/dL
55-<60 years: 3.87-14.7 ng/dL
60-<65 years: 3.67-13.9 ng/dL
65-<70 years: 3.47-13.0 ng/dL
70-<75 years: 3.28-12.2 ng/dL
75-<80 years: 3.08-11.3 ng/dL
80-<85 years: 2.88-10.5 ng/dL
85-<90 years: 2.69-9.61 ng/dL
90-<95 years: 2.49-8.76 ng/dL
95-100+ years: 2.29-7.91 ng/dL

Males (children):

<1 year: Term infants

1-15 days: 0.20-3.10 ng/dL*

16 days-1 year: Values decrease gradually from newborn (0.20-3.10 ng/dL) to prepubertal levels

*Forest MG, Cathiard AM, Bertrand JA. Total and unbound testosterone levels in the newborn and in normal and hypogonadal children: use of a sensitive radioimmunoassay for testosterone. J Clin Endocrinol Metab.

1973;36(6):1132-1142

1-8 years: <0.13 ng/dL

9 years: <0.13-0.45 ng/dL

10 years: <0.13-1.26 ng/dL

11 years: <0.13-5.52 ng/dL

12 years: <0.13-9.28 ng/dL

13 years: <0.13-12.6 ng/dL

14 years: 0.48-15.3 ng/dL

15 years: 1.62-17.7 ng/dL

16 years: 2.93-19.5 ng/dL

17 years: 4.28-20.9 ng/dL

18 years: 5.40-21.8 ng/dL

19 years: 5.36-21.2 ng/dL

Females (adult):

20-<25 years: <0.13-1.08 ng/dL

25-<30 years: <0.13-1.06 ng/dL

30-<35 years: <0.13-1.03 ng/dL

35-<40 years: <0.13-1.00 ng/dL

40-<45 years: <0.13-0.98 ng/dL
45-<50 years: <0.13-0.95 ng/dL
50-<55 years: <0.13-0.92 ng/dL
55-<60 years: <0.13-0.90 ng/dL
60-<65 years: <0.13-0.87 ng/dL
65-<70 years: <0.13-0.84 ng/dL
70-<75 years: <0.13-0.82 ng/dL
75-<80 years: <0.13-0.79 ng/dL
80-<85 years: <0.13-0.76 ng/dL
85-<90 years: <0.13-0.73 ng/dL
90-<95 years: <0.13-0.71 ng/dL
95-100+ years: <0.13-0.68 ng/dL

Females (children):

<1 year: Term infants

1-15 days: <0.13-0.25 ng/dL *

16 days-1 year: Values decrease gradually from newborn (<0.13-0.25 ng/dL) to prepubertal levels

*Forest MG, Cathiard AM, Bertrand JA. Total and unbound testosterone levels in the newborn and in normal and hypogonadal children: use of a sensitive radioimmunoassay for testosterone. J Clin Endocrinol Metab. 1973;36(6):1132-1142

1-4 years: <0.13 ng/dL
5 years: <0.13 ng/dL
6 years: <0.14 ng/dL
7 years: <0.13-0.23 ng/dL
8 years: <0.13-0.34 ng/dL
9 years: <0.13-0.46 ng/dL
10 years: <0.13-0.59 ng/dL
11 years: <0.13-0.72 ng/dL
12 years: <0.13-0.84 ng/dL
13 years: <0.13-0.96 ng/dL
14 years: <0.13-1.06 ng/dL
15-18 years: <0.13-1.09 ng/dL
19 years: <0.13-1.08 ng/dL

TESTOSTERONE, BIOAVAILABLE**Males**

< or =19 years: Not established

20-29 years: 83-257 ng/dL

30-39 years: 72-235 ng/dL

40-49 years: 61-213 ng/dL

50-59 years: 50-190 ng/dL

60-69 years: 40-168 ng/dL

> or =70 years: Not established

Females (non-oophorectomized)

< or =19 years: Not established

20-50 years (on oral estrogen): 0.80-4.0 ng/dL

20-50 years (not on oral estrogen): 0.80-10 ng/dL

>50 Years: Not established