

**Reporting Title:** Inhibin B, S**Performing Location:** Rochester**Ordering Guidance:**

For the initial evaluation of patients suspected of having a granulosa cell tumor of the ovary, order INHAB / Inhibin A and B, Tumor Marker, Serum. If the results of the profile show that either inhibin A or B are elevated, consider monitoring the patient with the individual tests, INHA / Inhibin A, Tumor Marker, Serum or INHB / Inhibin B, Serum.

**Specimen Requirements:**

Supplies: Sarstedt Aliquot Tube 5 mL (T914)

Collection Container/Tube:

Preferred: Serum gel

Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 0.5 mL

Collection Instructions: Centrifuge and aliquot serum into a plastic vial.

**Specimen Minimum Volume:**

0.4 mL

**Forms:**

If not ordering electronically, complete, print, and send an Oncology Test Request (T729) with the specimen.

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	14 days	
	Frozen	90 days	
	Ambient	7 days	

**Result Codes:**

Result ID	Reporting Name	Type	Unit	LOINC®
88722	Inhibin B, S	Numeric	pg/mL	56940-0

LOINC and CPT codes are provided by the performing laboratory.

**Supplemental Report:**

No

**CPT Code Information:**

83520

**Reference Values:**

## Males

<15 days: 68-373 pg/mL  
15-180 days: 42-516 pg/mL  
6 months-7 years: 24-300 pg/mL  
8-30 years: 47-383 pg/mL  
31-72 years: <358 pg/mL  
>72 years: Not established

## Females

< or =12 years: <183 pg/mL  
13-41 years Regular Cycle (Follicular Phase): <224 pg/mL  
42-51 years Regular Cycle (Follicular Phase): <108 pg/mL  
13-51 years Regular Cycle (Luteal Phase): <80 pg/mL  
>51 years (Postmenopausal): <12 pg/mL