

Reporting Title: C5-OH Acylcarnitine, QN, U**Performing Location:** Rochester**Ordering Guidance:**

This second-tier test is used specifically to evaluate a newborn screening elevation of 3-hydroxyisovaleryl-/2-methyl-3-hydroxy acylcarnitine and must not be ordered with either C4U / C4 Acylcarnitine, Quantitative, Random, Urine or C5DCU / C5-DC Acylcarnitine, Quantitative, Random, Urine.

For general screening for metabolic disorders, see OAU / Organic Acids Screen, Random, Urine; ACRN / Acylcarnitines, Quantitative, Plasma; and AAQP / Amino Acids, Quantitative, Plasma.

Necessary Information:

Include patient's age, family history, clinical condition (asymptomatic or acute episode), diet, and drug therapy information.

Specimen Requirements:

Patient Preparation: If clinically feasible, discontinue L-carnitine supplementation at least 72 hours before specimen collection.

Supplies: Urine Tubes, 10 mL (T068)

Collection Container/Tube: Clean, plastic urine collection container

Submission Container/Tube: Plastic, 10-mL urine tube

Specimen Volume: 5 mL

Collection Instructions:

1. Collect a random urine specimen.
2. Freeze specimen immediately.

Specimen Minimum Volume:

1 mL

Forms:

If not ordering electronically, complete, print, and send a Biochemical Genetics Test Request (T798) with the specimen.

Specimen Type	Temperature	Time	Special Container
Urine	Frozen (preferred)	7 days	
	Refrigerated	24 hours	

Result Codes:

Result ID	Reporting Name	Type	Unit	LOINC®
88830	C5-OH Acylcarnitine, QN, U	Numeric	mmol/mol Cr	50091-8
28125	C5-OH Interpretation	Alphanumeric		59462-2
34469	Reviewed By	Alphanumeric		18771-6

LOINC and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

82017

Reference Values:

<2.93 millimoles/mole creatinine