

Reporting Title: Antithrombin Activity, P**Performing Location:** Rochester**Ordering Guidance:**

Coagulation testing is highly complex, often requiring the performance of multiple assays and correlation with clinical information. For that reason, consider ordering AATHR / Thrombophilia Profile, Plasma and Whole Blood.

Specimen Requirements:

Specimen Type: Platelet-poor plasma

Collection Container/Tube: Light-blue top (3.2% sodium citrate)

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Collection Instructions:

1. For complete instructions, see Coagulation Guidelines for Specimen Handling and Processing.
2. Centrifuge, transfer all plasma into a plastic vial, and centrifuge plasma again.
3. Aliquot plasma into a plastic vial leaving 0.25 mL in the bottom of centrifuged vial.
4. Freeze plasma immediately (no longer than 4 hours after collection) at -20 degrees C or, ideally at -40 degrees C or below.

Additional Information:

1. A double-centrifuged specimen is critical for accurate results as platelet contamination may cause spurious results.
2. Each coagulation assay requested should have its own vial.
3. Heparin treatment may lower plasma antithrombin.

Specimen Minimum Volume:

0.5 mL

Forms:

If not ordering electronically, complete, print, and send a Coagulation Test Request (T753) with the specimen.

Specimen Type	Temperature	Time	Special Container
Plasma Na Cit	Frozen	14 days	

Result Codes:

Result ID	Reporting Name	Type	Unit	LOINC®
ATTF	Antithrombin Activity, P	Numeric	%	27811-9

LOINC and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

85300

Reference Values:

Normal values: 80-130%

Normal, full-term newborn infants have lower levels ($>$ or \approx 35-40%) that reach normal values by age 90 days.

Premature infants (30-36 weeks gestation) have lower levels that reach normal values by age 180 days.